



![](_page_1_Figure_1.jpeg)

![](_page_1_Picture_2.jpeg)

![](_page_2_Figure_1.jpeg)

![](_page_2_Figure_2.jpeg)

![](_page_2_Figure_3.jpeg)

![](_page_3_Figure_1.jpeg)

![](_page_3_Figure_2.jpeg)

![](_page_3_Picture_3.jpeg)

![](_page_4_Picture_1.jpeg)

Crisaborole's role on the Schachner Steroid Ladder: Topical Treatment Severity Schedule Clobetasel (high potency CS) Twice daily for 3-5 days + TCl of PDI + emollients If Severe: Triamcinolone (medium If Moderate: Twice daily for 3-5 days potency CS) + TCl or PDI emollients If Mild: Alclometasone (low potency Twice daily for 3-5 days CS) + TCI or (PDI)+ emollients TCI or (PDI) or TS + emollients Twice daily for 2 weeks Controlled: \* Antihistamines as needed TCI or PDI or TS + emollients Maintenance (to areas of Twice weekly for 6 months \* Antibiotics as needed predilection): Long-term Maintenance & Emollients Twice daily Prevention: \*Abbreviations: CS: Corticosteroid. PDI: Phosphodiesterase inhibitor. TCI: Topical Calcineurin Inhibitor Oberlin KE, Nanda S. Atopic dermatitis made easy: The Schachner Ladder. Pediatr Dermatol. 2019;36(6):1017-8.

![](_page_5_Figure_1.jpeg)

![](_page_5_Figure_3.jpeg)

![](_page_6_Figure_1.jpeg)

![](_page_6_Figure_3.jpeg)

![](_page_7_Figure_1.jpeg)

![](_page_7_Figure_2.jpeg)

![](_page_7_Picture_3.jpeg)

![](_page_8_Figure_1.jpeg)

![](_page_8_Picture_3.jpeg)

![](_page_9_Figure_1.jpeg)

![](_page_9_Figure_2.jpeg)

![](_page_10_Figure_1.jpeg)

![](_page_10_Figure_2.jpeg)

![](_page_11_Figure_1.jpeg)

![](_page_11_Figure_2.jpeg)

![](_page_11_Picture_3.jpeg)

![](_page_12_Picture_1.jpeg)

![](_page_12_Picture_3.jpeg)

![](_page_13_Picture_1.jpeg)

![](_page_13_Picture_3.jpeg)

![](_page_14_Figure_1.jpeg)

![](_page_14_Figure_2.jpeg)

![](_page_14_Picture_3.jpeg)

![](_page_15_Figure_1.jpeg)

![](_page_15_Figure_2.jpeg)

![](_page_15_Picture_3.jpeg)

![](_page_16_Figure_1.jpeg)

![](_page_16_Picture_3.jpeg)

![](_page_17_Figure_1.jpeg)

![](_page_17_Figure_2.jpeg)

![](_page_18_Figure_1.jpeg)

![](_page_18_Figure_2.jpeg)

![](_page_19_Picture_1.jpeg)

![](_page_19_Picture_3.jpeg)

![](_page_20_Figure_1.jpeg)

![](_page_20_Figure_3.jpeg)

![](_page_21_Figure_1.jpeg)

![](_page_21_Figure_2.jpeg)

![](_page_21_Picture_3.jpeg)

## Oral Anticholinergics: Adverse Effects May Impede Usage

- Efficacy is typically achieved at dosages that induce side effects (dry mouth, blurred vision, dry eyes, urinary retention)
- One-third of patients discontinue use due to side effects
- Gradually increasing the dosage may help reduce adverse effects
- Overheating when outdoors is of particular concern in children チョーマーマー

 Nawrocki S, Cha J. The etiology, diagnosis, and management of hyperhidrosis: A comprehensive review: Therapeutic options. J Am Acad Dermatol. 2019;81(3):669-80.
Gordon JR, Hill SE. Update on pediatric hyperhidrosis. Dermatol Ther. 2013;26(6):452-61.

![](_page_22_Figure_7.jpeg)

![](_page_22_Figure_8.jpeg)

![](_page_22_Picture_9.jpeg)

![](_page_23_Picture_1.jpeg)

![](_page_23_Picture_2.jpeg)

![](_page_23_Picture_3.jpeg)

![](_page_24_Figure_1.jpeg)